

# 論 說

## 臺灣植物分類雜纂 (6)

佐々木舜一

Miscellaneous contributions to the  
Flora of Formosa, VI.

by

Syun'iti SASAKI

---

### (38) タイワンボタソヅルの學名

本種は下記の學名を用ふるを至當と考へ次の如く訂正す。

*Clematis Gouriana* Roxb., Hort. Beng. (1814) 43, nomen nudum, Fl. Ind. II (1832) 670; DC., Syst. I (1818) 138, Prodr. I (1824) 3; Miquel, Fl. Ind. Batav. I. 2 (1859) 4; Hook. et Thoms., in Hook. Fl. Brit. Ind. I (1875) 4; Trim, Handb. Fl. Ceyl. I (1893) 2; Merrill, Enum. Philip. Fl. Pl. II (1923) 141; Sasaki, List Pl. Formosa (1928) 183.

*Clematis taiwaniana* Hayata, Kawakami, List Pl. Formos. (1910) 2; Mater. Fl. Formos. (1911) 17, Icon. Pl. Formos. I (1911) 23; Kawakami et Sasaki, in Trans. Natur. Hist. Soc. Formos. V. No. 22, ad. (1915) 3; Hayata, Gen. Ind. Fl. Formos. (1916) 2; Makino et Nemoto, Fl. Jap. (1925) 971.

Nom. Jap. Taiwan-botanduru. タイワンボタソヅル

Hab. Patina, Taihoku (T. Makino, 1896); Daihunpo, Taityū Pref. (T. Kawakami & U. Mori, Aug. 1904); Takusya, Taityū Pref. (U. Mori, July 1907); Mararaosya, Taitō (Z. Koboyasi, Aug. 1907); Taihoku (S. Sasaki, July 1909); Sōzan, Taihoku (S. Sasaki, Aug. 1910, July 1916); Sankakuyū, Tōyen (Y. Simada, Sep. 1913); Sōzan, Taihoku (Y. Simada, Aug. 1915); Hokuzankō, Nantō (Takeo Ito, Oct. 1916); Kōtō-syo (R. Kanehira & S. Sasaki, June 1919).

Distrib. India, Ceylon, Java and Philippine Isl.

Formos. (1916) ibid.; Makino et Nemoto, ibid.; Sasaki, ibid.

Nom. Jap. Taiwan-keman, Nantō-keman. タイワンケマン 又はナントウケマン

Hab. Hōran-zan, Taihoku (T. Sōma, March 1912); Uraisya, Taihoku (T. Kamiya & S. Sasaki, Feb. 1909); Kōkwaten, Tansui (Y. Simada, Apr. 1909); Mt. Sitisei (S. Sasaki, March 1910); Zyōsuisei, Tōyen (U. Mori, March 1907); Mt. Gosi, Sintiku (T. Kawakami & U. Mori, Jan. 1907); Mt. Ritō (T. Kawakami, July 1913); Mt. Nanko-taizan 6000-7000 ft. above the sea level (S. Sasaki, July 1922); Mt. Nōkō (E. Matuda, Aug. 1919); Taikwan, Niitaka 7000 ft. above the sea (S. Sasaki, Dec. 1923); Baikei, Nantō (R. Kanehira & S. Sasaki, March 1918); Sankakuyū (Sankkyō), Tōyen (T. Sōma, Apr. 1917); Mt. Niitaka 8000 ft. above the sea level (T. Kawakami & U. Mori, Oct. 1909); Mt. Arisan (T. Kawakami & U. Mori, March 1908); Karapin, Arisan 3000 ft. above the sea level (T. Sōma, Jan. 1915); Ahisen central mt. range, 4000 ft. above the sea level (T. Kawakami & S. Sasaki, Jan. 1911); Kōsyun (G. Nakahara, Dec. 1906); Daironkō-sya, Taitō (T. Kawakami & U. Mori, Nov. 1906); Taroko, Taitō (now Kwarenkō Pref.) (G. Nakahara, Jan. 1906); Hōka-syo (Agincourt) (T. Kawakami & U. Mori, May 1907); Taitō (S. Yasukawa, Feb. 1923).

Endemic plant.

#### (43) タイワンケマンの一變種

*Corydalis formosana* Hayata の一新變種を私等は本島山地に之を見ることが出來た、全株は殆んど基本種と變りはないが、葉の緣邊が悉く繊細なる缺刻に變化し、葉は基本種よりも小さい、そこで私はホソバキケマンの新和名を下し次の記載文を添加する。

*Corydalis formosana* Hayata, Mater. Fl. Formos. (1911) 26.

var. *microphylla* Sasaki, var. nov.

Annual or biennial glabrous herbs, reached 27-33 cm long, leaves fasciculate or scattered, caulis rather erect, peduncles or pedunculate. Leaves alternate bipinnate, thin, 8-11 cm long, pionae 10-20 mm long, leaflet 4 mm long, 4 mm wide, pointed, petioles 25-54 mm long, petiolules 3 mm long, serrate or incised of the margins. Flowers racemed, irregular, greenish yellow, 17 mm long; sepals small, petals exterior unequal, spure 4 mm long, 2mm broad, bracts obovate or elliptic, acute or acuminate of the apex, 8-9 mm long, 3-4 mm wide. Capsules linear, seeds orbicular compressed, minutely.

Nom. Jap. Hosoba-kikeman. ホソバキケマン (n.n.)

Hab. Takadai, 5000 feet above the sea level, Sintiku (S. Sasaki, March 1916)

Near *C. formosana* Hay., but differs from it in very thinner and minutely leaves.

(44) ホザキケマンの文献及其の產地

*Corydalis racemosa* Pers., Syn. Pl. II (1807) 270; DC., Syst. II (1821) 125, et Prodr. I (1843) 129; Miq., Prol. Fl. Jap. (1865) 200, pro parte; Franch. et Sav., Enum. Pl. Jap. I (1875) 30, pro parte, et II (1879) 275; Max., Pl. Chin. Potanin. in Act. Hort. Petrop. XI (1890) 49; Franch., Pl. David. I (1883) 30; Forbes et Hemsl., Ind. Fl. Sin. I (1886) 38; Henry, List Pl. Formos. (1896) 17; Ito et Matsum., Tent. Fl. Lutch. (1900) 28; Matsum. et Hayat., Enum. Pl. Formos. (1906) 21; Hayat., Icon. Pl. Formos. I (1911) 44, Gen. Ind. Fl. Formos. (1916) 4; Makino et Nemoto, Fl. Jap. (1925) 913; Sasaki, List Pl. Formos. (1928) 198.

*Fumaria racemosa* Willd., Sp. Pl. III<sup>2</sup> (1800) 864; Thunb., in "Nov. Act. Pterop. XII (1801) 103, t. b."

Nom. Jap. Hozaki-keman, Hozaki-kikeman, ホザキケマン又はホザキキケマン

Hab. Kiirun (S. Sasaki, Apr. 1910 & May 1923); Sekitei, Kiirun (S. Sasaki & S. Hudii, March 1913); Sekimon, Kiirun (S. Sasaki, Apr. 1910); Isl. Kiirun (S. Sasaki, May 1910); Kienpōri, Kiirun (S. Sasaki, Apr. 1910); Hūkikaku, near Tansui (T. Sōma, Nov. 1911 & S. Sasaki, March 1924); Hatune, Kwarenkō (E. Matuda, Aug. 1919); Ranrun, Nantō (Taityū-syū) (T. Kawakami & U. Mori, July 1907); Beiron, Taitō (Kwarenkō) (Z. Kabayasi, May 1906); Taitō (T. Sōma March 1913); Miyasan, Kwarenkō (S. Yasukawa, March 1923).

Distrib. Japan and China.

(45) コパンバヤナギ

本種は從來恒春半島の產とのみ知られたるも、其後臺中州蓮華池方面の森林にも多く自生するを知られた。而して早田博士は之に *Litosanthera gracilis* Hay. の學名を附して *Icones Plantarum Formosanarum* II に於て發表した、然るに一方非列賓に自生する *Litosanthera biflora* Blume の標本を Merrill 氏によりて寄贈を受け、之と本島產と對照するに全然同一種なることを明かにするを得たるを以て、私は茲に次の様に整理を行ひ、小著臺灣植物名彙を以て發表した。

*Litosanthera biflora* Blume, Cat. Gew. Buitenz. (1823) 21; DC., Prodr. IV (1830) 465; Merr., Enum. Philip. Fl. Pl. III (1923) 568; Sasaki, List Pl. Formos.