

Revisio ad Floram Formosanam. (I)

Takahide HOSOKAWA

1. *Satureia Kudoii* HOSOKAWA, *sp. nov.*

Herba suberecta, fere 30 cm. alta. Caulis rigidiusculus, basi patentim puberus, sursum patentim hirtellus. Folia petiolata, rhombeo-ovata, apice obtusa, basi cuneata, margine crenato-serrata, chartaceo-membrancea, supra viridia, scabra, hirtello-pubescentia, subtus paridiora, glandulosa, pilis sparsis hirsutiuscula, costis venisque elevatis, ad 3 cm. longa, ad 2 cm. lata, petiolo 0.6-1.5 cm. longo dense hirsuto. Inflorescentia terminalis, rhachide villosa-hirsuto; verticillastri densiflori, in spicam densam approximati, pedicellis hirtellis 2 mm. longis, bracteolis linearibus margine longe pilosis ad 2.5 mm. longis. Calyx subrecurvatus, tubulosus, pilis longis mollibusque hirsutus, dorso apiceque purpureus, bilabiatus, ad 5 mm. longus, labio superiore 3-dentato, dentibus subulato-deltoides, labio inferiore 2-fido, fidis subulato-linearibus. Corolla 8 mm. longa, extus pubescens, labio superiore rotundato apice emarginato, inferiore patente 3-lobo, lobo medio subrotundato.

NON. JAP. *Taiwan-miyamatōbana* (*nov.*).

HAB. Inter Sikikun et Piyanan-ambu, Praef. Taihoku (IPSE, n. 2389! Jul. 10, 1931; Typus in Herb. Univ. Imp. Taihoku).

Arcte affinis *S. chinensi*, sed inflorescentiis terminalibus dense spicatis, et foliis subtus glanduliferis, ex qua differt.

2. *Carpesium spathiforme* HOSOKAWA, *sp. nov.*

Suffrutex circ. 70 cm. altus. Caulis erectus, subcanaliculatus, adpresse pilosus. Folia radicalia numquam rosulata, elliptico-spathulata, apice obtusa, basi in petiolum alatum sensim attenuatissima, margine integra, membranacea, utraque pilosa, sed costis subtus pilis longis adpressisque sericeis, saepius in maximum 12 cm. longa, 4 cm. lata. Inflorescentia subcorymboso-panicu-